

S-1+NDP 療法

適応	子宮頸癌	催吐性リスク	中等度
----	------	--------	-----

			Day1	Day8	Day15	Day22	Day29	Day35
S-1	40mg/m ² /回	1日2回, 経口	→			21日間		
NDP	90mg/m ²	DIV, 90min		●				

5週間毎, PDまで

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35			
① S-1 40mg/m ² ×2		◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎																	
② グラニセトロンバッグ3mg 50mL									●																														
デキサート注射液 9.9mg									●																														
DIV 30min																																							
③ アクブラ 90mg/m ²									●																														
生理食塩液注 500mL									●																														
DIV 90min																																							
④ ソルアセトF輸液 500mL									●																														
DIV 90min																																							
⑤ ソルデム3A輸液 500mL									●																														
DIV 90min																																							

【S-1投与量】

体表面積	S-1初回基準量
1.25m ² 未満	40mg/回
1.25m ² 以上1.5m ² 未満	50mg/回
1.5m ² 以上	60mg/回