

# Pembrolizumab+Axitinib 療法

|    |      |            |     |
|----|------|------------|-----|
| 適応 | 腎細胞癌 | 催吐性<br>リスク | 最小度 |
|----|------|------------|-----|

| 200           |            |            | Day1   | Day8 | Day15 | Day21 |
|---------------|------------|------------|--------|------|-------|-------|
| Axitinib      | 5mg/回      | 1日2回, 経口   | —————▶ |      |       |       |
| Pembrolizumab | 200mg/body | DIV, 30min | ●      |      |       |       |

3週間毎, PDまで

| 400           |            |            | Day1   | Day8 | Day15 | Day22 | Day29 | Day36 | Day42 |
|---------------|------------|------------|--------|------|-------|-------|-------|-------|-------|
| Axitinib      | 5mg/回      | 1日2回, 経口   | —————▶ |      |       |       |       |       |       |
| Pembrolizumab | 400mg/body | DIV, 30min | ●      |      |       |       |       |       |       |

6週間毎, PDまで

| 200 |                  | Day | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
|-----|------------------|-----|---|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| ①   | インライタ 5mg/回×2    |     |   | ○ | ○ | ○ | ○ | ○ | ○ | ○ | ○ | ○ | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○  |    |    |    |    |    |    |    |    |
|     | 経口               |     |   |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ②   | キトラーダ 200mg/body |     |   | ● |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
|     | 生理食塩液注 100mL     |     |   | ● |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
|     | DIV 30min        |     |   |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ③   | 生理食塩液注 50mL      |     |   | ● |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
|     | DIV 15min        |     |   |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |

  

| 400 |                  | Day | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | ***** |       |       |       |       |       | 42 |
|-----|------------------|-----|---|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|-------|-------|-------|-------|-------|-------|----|
| ①   | インライタ 5mg/回×2    |     |   | ○ | ○ | ○ | ○ | ○ | ○ | ○ | ○ | ○ | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○  | ○     | ***** | ***** | ***** | ***** | ***** | ○  |
|     | 経口               |     |   |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    | ***** | ***** | ***** | ***** | ***** |       |    |
| ②   | キトラーダ 400mg/body |     |   | ● |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    | ***** | ***** | ***** | ***** | ***** |       |    |
|     | 生理食塩液注 100mL     |     |   | ● |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    | ***** | ***** | ***** | ***** | ***** |       |    |
|     | DIV 30min        |     |   |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    | ***** | ***** | ***** | ***** | ***** |       |    |
| ③   | 生理食塩液注 50mL      |     |   | ● |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    | ***** | ***** | ***** | ***** | ***** |       |    |
|     | DIV 15min        |     |   |   |   |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    | ***** | ***** | ***** | ***** | ***** |       |    |

【Avelumab前投薬】 レスタミンコーワ錠 50mg, カロナール錠 600mg/経口