

SSP (S-1+CDDP分割) 療法

適応	胃癌	催吐性リスク	高度
----	----	--------	----

			Day1	Day8	Day15	Day22	Day29	Day35	
S-1	40mg/m ² /回	1日2回, 経口	→ 21日間						
CDDP	30mg/m ²	DIV, 60min	●		●				

5週間毎, PDまで

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	
① S-1 40mg/m ² ×2		○	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	○														
経口		夕																				朝															
② ビカネイト輸液 1,000mL		●														●																					
DIV 300min																																					
③ マンニトールS注射液 300mL		●														●																					
DIV 45min																																					
④ アロキシバッグ0.75mg 50mL		●														●																					
デキサート注射液 9.9mg		●														●																					
DIV 15min																																					
⑤ シスプラチン 30mg/m ²		●														●																					
生理食塩液注 250mL		●														●																					
DIV 60min																																					

【S-1投与量】

体表面積	S-1初回基準量
1.25m ² 未満	40mg/回
1.25m ² 以上1.5m ² 未満	50mg/回
1.5m ² 以上	60mg/回