

XELOX+Bev 療法

適応

大腸癌

催吐性
リスク

中等度

			Day1	Day8	Day15	Day21	
Capecitabine	1,000mg/m ² /回	1日2回, 経口	—————→			14日間	
Bevacizumab	7.5mg/kg	DIV, 90→60→30min	●				
L-OHP	130mg/m ²	DIV, 120min	●				

3週間毎,

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① カペシタピン 1000mg/m ² ×2		○	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	○													
経口		夕														朝													
② アロキシバグ0.75mg	50mL	●																											
デキサート注射液	9.9mg	●																											
	DIV 15min																												
③ アバスチン 7.5mg/kg		●																											
生理食塩液注	100mL	●																											
	DIV 90→60→30min																												
④ オキサリプラチン 130mg/m ²		●																											
5%ブドウ糖液	500mL	●																											
	DIV 120min																												

【Capecitabine投与量】

体表面積	1回用量
1.36m ² 未満	1,200mg/回
1.36m ² 以上1.66m ² 未満	1,500mg/回
1.66m ² 以上1.96m ² 未満	1,800mg/回
1.96m ² 以上	2,100mg/回