

# GCS (GEM+CDDP+S-1) 療法

適応	胆道癌	催吐性 リスク	高度
----	-----	------------	----

			Day1	Day8	Day14
S-1	40mg/m <sup>2</sup> /回	1日2回, 経口	→	7日間	
GEM	1,000mg/m <sup>2</sup>	DIV, 30min	●		
CDDP	25mg/m <sup>2</sup>	DIV, 60min	●		

2週間毎, PDまで

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① S-1 40mg/m <sup>2</sup> ×2		○	◎	◎	◎	◎	◎	◎	○																				
経口		夕							朝																				
② ビカネイト輸液 1,000mL		●							●																				
DIV 300min																													
③ マンニトールS注射液 300mL		●							●																				
DIV 45min																													
④ アロキシバッグ0.75mg 50mL		●							●																				
デキサート注射液 9.9mg																													
DIV 15min																													
⑤ ゲムシタピン 1,000mg/m <sup>2</sup>		●							●																				
生理食塩液注 100mL		●							●																				
DIV 30min																													
⑥ シスプラチン 25mg/m <sup>2</sup>		●							●																				
生理食塩液注 250mL		●							●																				
DIV 60min																													
⑥ 生理食塩液注 50mL		●							●																				
DIV 15min																													

【S-1投与量】

体表面積	S-1初回基準量
1.25m <sup>2</sup> 未満	40mg/回
1.25m <sup>2</sup> 以上1.5m <sup>2</sup> 未満	50mg/回
1.5m <sup>2</sup> 以上	60mg/回