

TP+Bev 療法

適応	子宮頸癌	催吐性 リスク	高度
----	------	------------	----

			Day1	Day8	Day15	Day21
PTX	175mg/m ²	DIV, 180min	●			
CDDP	50mg/m ²	DIV, 60min	●			
Bevacizumab	15mg/kg	DIV, 90→60→30min	●			

3週間毎, 6コース

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① ソルデム1輸液	500mL	●																											
硫酸Mg補正液1mEq/mL	8mL	●																											
	DIV 60min																												
② 生理食塩液注	50mL	●																											
デキサート注射液	9.9mg	●																											
ファモチジン注	20mg	●																											
	DIV 15min																												
③ バクリタキセル 175mg/m ²		●																											
生理食塩液注	500mL	●																											
	DIV 180min																												
④ アロキシバグ0.75mg	50mL	●																											
	DIV 15min																												
⑤ マンニトールS注射液	300mL	●																											
	DIV 30min																												
⑥ シスプラチン 50mg/m ²		●																											
生理食塩液注	500mL	●																											
	DIV 60min																												
⑦ ソルデム3A輸液	500mL	●																											
	DIV 60min																												
⑧ アバスチン 15mg/kg		●																											
生理食塩液注	100mL	●																											
	DIV 90→60→30min																												

【PTX前投薬】レスタミンコーワ錠 50mg/経口