

Pembrolizumab 200/400 療法

| | | | |
|----|-----------------|------------|-----|
| 適応 | MSI-High 固形癌 | 催吐性 リスク | 最小度 |
|----|-----------------|------------|-----|

| 200 | | | Day1 | Day8 | Day15 | Day21 |
|---------------|------------|------------|------|------|-------|-------|
| Pembrolizumab | 200mg/body | DIV, 30min | ● | | | |

3週間毎, PDまで

| 400 | | | Day1 | Day8 | Day15 | Day22 | Day29 | Day36 | Day42 |
|---------------|------------|------------|------|------|-------|-------|-------|-------|-------|
| Pembrolizumab | 400mg/body | DIV, 30min | ● | | | | | | |

6週間毎, PDまで

200

| Day | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
|---------------------|---|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| ① キイトルーダ 200mg/Body | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 生理食塩液注 100mL | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIV 30min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ② 生理食塩液注 50mL | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIV 15min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

400

| Day | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | ***** | 42 |
|---------------------|---|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|-------|----|
| ① キイトルーダ 400mg/Body | | ● | | | | | | | | | | | | | | | | | | | | | ***** | |
| 生理食塩液注 100mL | | ● | | | | | | | | | | | | | | | | | | | | | ***** | |
| DIV 30min | | | | | | | | | | | | | | | | | | | | | | | ***** | |
| ② 生理食塩液注 50mL | | ● | | | | | | | | | | | | | | | | | | | | | ***** | |
| DIV 15min | | | | | | | | | | | | | | | | | | | | | | | ***** | |