

CDDP+ALIMTA 療法

適応

非小細胞肺癌

催吐性
リスク

高度

			Day1	Day8	Day15	Day21
CDDP	75mg/m ²	DIV, 60min	●			
PEM	500mg/m ²	DIV, 10min	●			

3週間毎, 4コース

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① ソルデム1輸液	500mL	●																											
硫酸Mg補正液1mEq/mL	8mL	●																											
DIV	60min																												
② アロキシバッグ0.75mg	50mL	●																											
デキサート注射液	9.9mg	●																											
DIV	30min																												
③ アリムタ 500mg/m ²		●																											
生理食塩液注	100mL	●																											
DIV	10min																												
④ マンニトールS注射液	300mL	●																											
DIV	30min																												
⑤ シスプラチン 75mg/m ²		●																											
生理食塩液注	500mL	●																											
DIV	60min																												
⑥ ソルデム3A輸液	500mL	●																											
DIV	60min																												

【初回投与7日前から開始】パンビタン末：1.0g/回，経口、連日

【初回投与7日前】メチコバル注：1,000μg/回、筋注，9週毎

※最終投与日から22日目まで継続