

CDDP+PEM+BV 療法

適応

非小細胞肺癌

催吐性
リスク

高度

			Day1	Day8	Day15	Day21
CDDP	75mg/m ²	DIV, 60min	●			
PEM	500mg/m ²	DIV, 10min	●			
Bevacizumab	15mg/kg	DIV, 90→60→30min	●			

3週間毎, 4コース

	Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① ソルデム1輸液	500mL		●																											
硫酸Mg補正液1mEq/mL	8mL		●																											
	DIV 60min																													
② アロキシバッグ0.75mg	50mL		●																											
デキサート注射液	9.9mg		●																											
	DIV 30min																													
③ アリムタ 500mg/m ²			●																											
生理食塩液注	100mL		●																											
	DIV 10min																													
④ マンニトールS注射液	300mL		●																											
	DIV 30min																													
⑤ シスプラチン 75mg/m ²			●																											
生理食塩液注	500mL		●																											
	DIV 60min																													
⑥ アバスチン 15mg/kg			●																											
生理食塩液注	100mL		●																											
	DIV 90→60→30min																													
⑥ ソルデム3A輸液	500mL		●																											
	DIV 60min																													

【初回投与7日前から開始】パンビタン末：1.0g/回，経口、連日， 【初回投与7日前】メチコバル注：1,000μg/回、筋注，9週毎

※最終投与日から22日目まで継続