

Pembro+CBDCA+PEM 療法

適応

非小細胞肺癌

催吐性
リスク

中等度*

			Day1	Day8	Day15	Day21
Pembrolizumab	200mg/body	DIV, 30min	●			
PEM	500mg/m ²	DIV, 10min	●			
CBDCA	AUC 5	DIV, 60min	●			

3週間毎, 4-6コース

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① キイトルーダ 200mg/body		●																											
生理食塩液注	100mL	●																											
DIV	30min																												
② 生理食塩液注	50mL	●																											
DIV	15min																												
③ グラニセトロンバッグ3mg	50mL	●																											
デキサート注射液	4.95mg	●																											
DIV	30min																												
④ アリムタ 500mg/m ²		●																											
生理食塩液注	100mL	●																											
DIV	10min																												
⑤ カルボプラチン AUC 5		●																											
5%ブドウ糖液	250mL	●																											
DIV	60min																												

【初回投与7日前から開始】パンピタン末：1.0g/回、経口、連日

【初回投与7日前】メチコバル注：1,000μg/回、筋注、9週毎

※最終投与日から22日目まで継続