

Atezo+Bev+CP 療法

適応

非小細胞肺癌

催吐性
リスク

中等度*

| | | | Day1 | Day8 | Day15 | Day21 |
|--------------|----------------------|------------------|------|------|-------|-------|
| Atezolizumab | 1,200mg/body | DIV, 60→30min | ● | | | |
| Bevacizumab | 15mg/kg | DIV, 90→60→30min | ● | | | |
| PTX | 200mg/m ² | DIV, 180min | ● | | | |
| CBDCA | AUC 6 | DIV, 60min | ● | | | |

3週間毎, 4コース

| Day | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
|--------------------------------|---|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| ① テセントリク 1,200mg/body | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 生理食塩液注 250mL | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIV 60→30min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ② アバスチン 15mg/kg | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 生理食塩液注 100mL | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIV 90→60→30min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ③ 生理食塩液注 50mL | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| デキサート注射液 19.8mg | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ファモチジン注 20mg | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIV 15min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ④ グラニセトロンバッグ3mg 50mL | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIV 30min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ⑤ パクリタキセル 200mg/m ² | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 生理食塩液注 500mL | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIV 180min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ⑥ カルボプラチン AUC 6 | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 5%ブドウ糖液 250mL | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| DIV 60min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

【PTX前投薬】 レスタミンコーワ錠 50mg/経口