

# weekly PTX+Trastuzumab 療法

|    |    |            |    |
|----|----|------------|----|
| 適応 | 乳癌 | 催吐性<br>リスク | 軽度 |
|----|----|------------|----|

|             |                     |               | Day1 | Day8 | Day15 | Day22 | Day28 |
|-------------|---------------------|---------------|------|------|-------|-------|-------|
| Trastuzumab | 4mg→2mg/kg          | DIV, 90→30min | ●    | ●    | ●     | ●     |       |
| PTX         | 80mg/m <sup>2</sup> | DIV, 60min    | ●    | ●    | ●     |       |       |

4週間毎, PDまで

| Day                           | 0 | 1 | 2         | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
|-------------------------------|---|---|-----------|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| ① トラスツズマブ 4mg→2mg/kg          |   | ● |           |   |   |   |   |   | ● |   |    |    |    |    |    | ●  |    |    |    |    |    |    | ●  |    |    |    |    |    |    |
| 蒸留水                           |   | ● |           |   |   |   |   |   | ● |   |    |    |    |    |    | ●  |    |    |    |    |    |    | ●  |    |    |    |    |    |    |
| 生理食塩液注 250mL                  |   | ● |           |   |   |   |   |   | ● |   |    |    |    |    |    | ●  |    |    |    |    |    |    | ●  |    |    |    |    |    |    |
| DIV 90→30min                  |   |   |           |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ② 生理食塩液注 50mL                 |   | ● |           |   |   |   |   |   | ● |   |    |    |    |    |    | ●  |    |    |    |    |    |    |    |    |    |    |    |    |    |
| デキサート注射液 6.6mg                |   | ● | 初回は8.25mg |   |   |   |   |   |   | ● |    |    |    |    | ●  |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ファモチジン注 20mg                  |   | ● |           |   |   |   |   |   | ● |   |    |    |    |    |    | ●  |    |    |    |    |    |    |    |    |    |    |    |    |    |
| DIV 15min                     |   |   |           |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ③ 生理食塩液注 50mL                 |   | ● |           |   |   |   |   |   | ● |   |    |    |    |    |    | ●  |    |    |    |    |    |    |    |    |    |    |    |    |    |
| DIV 30min                     |   |   |           |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
| ④ パクリタキセル 80mg/m <sup>2</sup> |   | ● |           |   |   |   |   |   | ● |   |    |    |    |    |    | ●  |    |    |    |    |    |    |    |    |    |    |    |    |    |
| 生理食塩液注 250mL                  |   | ● |           |   |   |   |   |   | ● |   |    |    |    |    |    | ●  |    |    |    |    |    |    |    |    |    |    |    |    |    |
| DIV 60min                     |   |   |           |   |   |   |   |   |   |   |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |

【PTX前投薬】 レスタミンコーワ錠 50mg/経口