

HP+PTX 療法

適応	乳癌	催吐性 リスク	軽度
----	----	------------	----

			Day1	Day8	Day15	Day21
Pertuzumab	840mg→420mg/body	DIV, 60→30min	●			
Trastuzumab	8mg→6mg/kg	DIV, 90→30min	●			
PTX	80mg/m ²	DIV, 60min	●	●	●	

3週間毎, PDまで

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① パーージェタ 840mg→420mg/body		●																											
生理食塩液注 250mL		●																											
DIV 60→30min																													
② 生理食塩液注 100→50mL		●																											
DIV 60→30min																													
③ トラスツズマブ 8mg→6mg/kg		●																											
蒸留水		●																											
生理食塩液注 250mL		●																											
DIV 90→30min																													
④ 生理食塩液注 50mL		●							●							●													
デキサート注射液 6.6mg		●	初回は8.25mg						●							●													
ファモチジン注 20mg		●							●							●													
DIV 15min																													
⑤ 生理食塩液注 50mL		●							●							●													
DIV 30min																													
⑥ バクリタキセル 80mg/m ²		●							●							●													
生理食塩液注 250mL		●							●							●													
DIV 60min																													

【前投薬】 レスタミンコーワ錠 50mg/経口